arXiv:1603.00650v1 [math.GN] 2 Mar 2016

LIMITS OF SEQUENCES OF CONTINUOUS FUNCTIONS DEPENDING ON FINITELY MANY COORDINATES

OLENA KARLOVA AND VOLODYMYR MYKHAYLYUK

Abstract. We answer two questions from *V.Bykov, On Baire class one functions on a product space, Topol. Appl. 199 (2016) 55–62,* and prove that every Baire one function on a subspace of a countable perfectly normal product is the pointwise limit of a sequence of continuous functions, each depending on finitely many coordinates. It is proved also that a lower semicontinuous function on a subspace of a countable perfectly normal product is the pointwise limit of an increasing sequence of continuous functions, each depending on finitely many coordinates, if and only if the function has a minorant which depends on finitely many coordinates.

1. Introduction

The collection of all continuous maps between topological spaces X and Y is denoted by $\mathrm{C}(X,Y)$. If $A \subseteq Y^X$, then we denote the family of all pointwise limits of sequences of maps from A by $\overline{A}^{\mathrm{p}}$. We say that a subset A of a topological space X is *functionally F_σ (G_δ)*, if it is a union (an intersection) of a sequence of functionally closed (functionally open) sets in X.

Recall that a map $f : X \to Y$ belongs to the *first Lebesgue class*, if the preimage $f^{-1}(V)$ of each open set $V \subseteq Y$ is an F_σ-set in X, and belongs to the *first Baire class*, if $f \in \overline{\mathrm{C}(X,Y)}^{\mathrm{p}}$. We denote the collection of all Lebesgue one maps and Baire one maps between X and Y by $\mathrm{H}_1(X,Y)$ and $\mathrm{B}_1(X,Y)$, respectively. If a Lebesgue (Baire) one map $f : X \to Y$ takes finitely many values, then we say that $f \in \mathrm{H}_1^0(X,Y)$ ($\mathrm{B}_1^0(X,Y)$). Let us observe that $\mathrm{H}_1(X,\mathbb{R}) = \mathrm{B}_1(X,\mathbb{R})$ for a perfectly normal space X (see [7, §31]).

Properties of functions defined on products of topological spaces are tightly connected with the dependence of such functions on some sets of coordinates. The dependence of continuous mappings on some sets of coordinates was investigated by many mathematicians (see, for instance [8], [2], [10] and the literature given there). However, the dependence on countably many coordinates serves as a convenient tool for the investigation of Namioka property of separately continuous maps and their analogs [9]. Functions depending locally on finitely many coordinates plays an important role in the theory of smoothness and renorming on Banach spaces [3].

Let $(X_n)_{n=1}^\infty$ be a sequence of topological spaces, $P = \prod_{n=1}^\infty X_n$ and $a = (a_n)_{n\in\mathbb{N}} \in P$ be a point. For every $n \in \mathbb{N}$ and $x \in P$ we put $p_n(x) = (x_1, \dots, x_n, a_{n+1}, a_{n+2}, \dots)$. We say that a set $A \subseteq P$ *depends on finitely many coordinates*, if there exists $n \in \mathbb{N}$ such that for all $x \in A$ and $y \in P$ the equality $p_n(x) = p_n(y)$ implies $y \in A$. A map $f : X \to Y$ defined on a subspace $X \subseteq P$ is *finitely determined* if it *depends on finitely many coordinates*, i.e., $f(x) = f(y)$ for all $x, y \in X$ with $p_n(x) = p_n(y)$. We denote by $\mathrm{CF}(X,Y)$ the set of all continuous finitely determined maps between X and Y; we write $\mathrm{CF}(X)$ for $\mathrm{CF}(X,\mathbb{R})$.

Vladimir Bykov in his recent paper [1] proved the following results.

Theorem A. *Let X be a subspace of a product $P = \prod_{n=1}^\infty X_n$ of a sequence of metric spaces X_n. Then*

1. *every Baire class one function $f : X \to \mathbb{R}$ is the pointwise limit of a sequence of functions in $\mathrm{CF}(X)$;*
2. *a lower semicontinuous function $f : X \to \mathbb{R}$ is the pointwise limit of an increasing sequence of functions from $\mathrm{CF}(X)$ if and only if f has a minorant in $\mathrm{CF}(X)$.*

The following questions were formulated in [1]:

Question 1. [1, Remark 1] *Does the conclusion (1) of Theorem A holds for an arbitrary completely regular space X?*

Question 2. [1, Remark 2] *Does the conclusion (2) of Theorem A holds for an arbitrary perfectly normal space X?*

We show that that the answer on Question 1 is negative (see Theorem 16). We prove that every Lebesgue one function $f : X \to Y$ is a pointwise limit of a sequence of functions from $\mathrm{CF}(X, Y)$ if Y is a path-connected separable metric R-space (in particular, a convex subset of a normed separable space) and one of the following properties holds: X is a Lindelöf subspace of a completely regular product P, or X is a subspace of perfectly normal P (see Theorem 8), or $X = P$ and P is pseudocompact (see Theorem 13). Theorem 9 gives the positive answer on Question 2.

2. Approximation of functions with finitely many values

Let X be a topological space. For a countable family $\mathcal{F} = (f_n : n \in \mathbb{N})$ of continuous functions $f_n : X \to [0, 1]$ we put

$$d_{\mathcal{F}}(x, y) = \sum_{n=1}^{\infty} \frac{1}{2^n} |f_n(x) - f_n(y)| \tag{2.1}$$

for all $x, y \in X$. Then $d_{\mathcal{F}}$ is a pseudometric on X such that every continuous function $f : (X, d_{\mathcal{F}}) \to Y$ is continuous on X in the initial topology for any topological space Y.

Lemma 1. *Let $P = \prod_{n=1}^{\infty} X_n$ be a product of completely regular spaces, X be a subspace of P and G be a functionally open set in X. If one of the following conditions holds:*

1. *P is perfectly normal, or*
2. *X is Lindelöf,*

then

(a) *there exists an increasing sequence of functions $f_n \in \mathrm{CF}(P, [0, 1])$ such that the characteristic function $\chi_G : X \to [0, 1]$ is a pointwise limit of the sequence $(f_n|_X)_{n=1}^{\infty}$;*

(b) *there exists a countable family $\mathcal{F}$ of functions from $\mathrm{CF}(P, [0, 1])$ such that G is open in $(X, d_{\mathcal{F}})$.*

Proof. **(a)**. Let $\tilde{G}$ be an open set in P such that $G = X \cap \tilde{G}$. Take a family $(W_s : s \in S)$ of basic functionally open sets in P such that

$$\tilde{G} = \bigcup_{s \in S} W_s.$$

Notice that G is an F_σ-set in X. Consequently, G is Lindelöf in case (2) and we choose at most countable set $S_0 \subseteq S$ such that $G \subseteq \bigcup_{s \in S_0} W_s$. Let us observe that $G = (\bigcup_{s \in S_0} W_s) \cap X$.

Denote $T = S$ in case (1) and $T = S_0$ in case (2). For every $k \in \mathbb{N}$ we put

$$T_k = \{s \in T : W_s \text{ depends on the first } k \text{ coordinates}\},$$

$$U_k = \bigcup_{s \in T_k} W_s.$$

Then U_k depends on the first k coordinates and $U_k \subseteq U_{k+1}$ for every $k \in \mathbb{N}$. Since for every $k \in \mathbb{N}$ the set U_k is functionally open in P, there exist sequences $(F_{k,n})_{n=1}^{\infty}$ and $(G_{k,n})_{n=1}^{\infty}$ of functionally closed and functionally open subsets of P, respectively, such that each of these sets depends on the first k coordinates,

$$U_k = \bigcup_{n=1}^{\infty} F_{k,n} \quad \text{and} \quad F_{k,n} \subseteq G_{k,n} \subseteq F_{k,n+1}.$$

For all $n \in \mathbb{N}$ we set

$$A_n = \bigcup_{k=1}^{n} F_{k,n} \quad \text{and} \quad B_n = \bigcup_{k=1}^{n} G_{k,n}.$$

Then the set A_n is functionally closed, the set B_n is functionally open, each of them depends on finitely many coordinates and $A_n \subseteq B_n \subseteq A_{n+1}$ for every n. Moreover, $G = (\bigcup_{n=1}^{\infty} A_n) \cap X$. We choose a function $f_n \in \mathrm{CF}(P, [0,1])$ such that

$$A_n = f_n^{-1}(1) \quad \text{and} \quad P \setminus B_n = f_n^{-1}(0)$$

for every n. Then $f_n(x) \le f_{n+1}(x)$ for all $n \in \mathbb{N}$ and $x \in P$. Moreover, $\lim_{n\to\infty} f_n(x) = \chi_G(x)$ for every $x \in X$.

(b). We put $\mathcal{F} = (f_n : n \in \mathbb{N})$, where $(f_n)_{n=1}^{\infty}$ is the sequence constructed in (a). Since each function f_n is continuous on $(P, d_{\mathcal{F}})$, the function χ_G is lower semicontinuous on $(X, d_{\mathcal{F}})$. Therefore, G is open in $(X, d_{\mathcal{F}})$. □

Proposition 2. *Let $P = \prod_{n=1}^{\infty} X_n$ be a product of completely regular spaces, X be a subspace of P and A be a functionally F_σ- and functionally G_δ-set in X simultaneously. If one of the following conditions holds:*

(1) *P is perfectly normal, or*
(2) *X is Lindelöf,*

then $\chi_A \in \overline{\mathrm{CF}(X, [0,1])}^{\mathrm{p}}$.

Proof. Lemma 1 implies the existence of a countable family $\mathcal{F}$ of functions from $\mathrm{CF}(P, [0,1])$ such that A and $X \setminus A$ are F_σ in $(X, d_{\mathcal{F}})$. Applying [1, Lemma 3] we obtain increasing sequences $(F_k^i)_{k=1}^{\infty}$ of closed sets in $(X, d_{\mathcal{F}})$ for $i = 1, 2$, satisfying the conditions

$$A = \bigcup_{k=1}^{\infty} F_k^1, \quad X \setminus A = \bigcup_{k=1}^{\infty} F_k^2,$$
$$d_k = d_{\mathcal{F}}(F_k^1, F_k^2) > 0 \quad \forall k \in \mathbb{N}.$$

For every $k \in \mathbb{N}$ we define a function $h_k : P \to [0,1]$ by the formula

$$h_k(x) = \min\{\tfrac{1}{d_k} d_{\mathcal{F}}(x, F_k^2), 1\}.$$

Now for every $k \in \mathbb{N}$ we put

$$f_k = h_k \circ p_k$$

and show that $(f_k)_{k=1}^{\infty}$ converges to χ_A pointwisely on X. Fix $x \in X$. Since

$$\lim_{k\to\infty} |f_k(x) - \chi_A(x)| = \lim_{k\to\infty} |h_k(p_k(x)) - h_k(x)| \le \lim_{k\to\infty} \frac{d_{\mathcal{F}}(p_k(x), x)}{d_k} = 0,$$

we have

$$\lim_{k\to\infty} f_k(x) = \chi_A(x).$$

The continuity of f_k on $(X, d_{\mathcal{F}})$ implies the continuity of f_k on X with the product topology induced from P. Hence, $\chi_A \in \overline{\mathrm{CF}(X)}^{\mathrm{p}}$. □

Remark 1. *It follows from* [6, Proposition 2.6] *that for every map $f \in \mathrm{H}_1(X, Y)$ the preimage $f^{-1}(V)$ of any open set $V \subseteq Y$ is functionally F_σ in X, if X is a normal space and Y is a metrizable space.*

Corollary 3. *Let P and X be as in Proposition 2. Then $\mathrm{H}_1^0(X) \subseteq \overline{\mathrm{CF}(X)}^{\mathrm{p}}$.*

Proof. We consider a function $f \in \mathrm{H}_1^0(X)$ with a finite image $f(X) = \{y_1, \dots, y_n\}$. Notice that the family $(A_i : i \in \{1, \dots, n\})$ of the sets $A_i = f^{-1}(y_i)$ forms a partition of X by functionally F_σ-sets. Moreover, $f(x) = \sum_{i=1}^{n} y_i \cdot \chi_{A_i}(x)$ for every $x \in X$. Since $y_i \cdot \chi_{A_i} \in \overline{\mathrm{CF}(X)}^{\mathrm{p}}$ for every i by Proposition 2, $f \in \overline{\mathrm{CF}(X)}^{\mathrm{p}}$. □

3. R-spaces and uniform limits

For a family $(f_s : s \in S)$ of maps between topological spaces X and Y we define a map $\Delta_{s\in S} f_s : X \to Y^S$ as the following:

$$\underset{s\in S}{\Delta} f_s(x) = (f_s(x))_{s\in S}$$

for every $x \in X$. We say that $\mathcal{F}$ is Δ-*closed*, if

$$f \circ (\underset{s\in S}{\Delta} f_s) \in \mathcal{F}$$

for any finite set $\{f_s : s \in S\} \subseteq \mathcal{F}$ and a continuous map $f : Y^S \to Y$.

A metric space (Y, d) is said to be an *R-space* [5], if for every $\varepsilon > 0$ there exists a continuous map $r_\varepsilon : Y \times Y \to Y$ with the following properties:

$$d(y, z) \le \varepsilon \implies r_\varepsilon(y, z) = y, \tag{3.1}$$

$$d(r_\varepsilon(y, z), z) \le \varepsilon \tag{3.2}$$

for all $y, z \in Y$. Let us observe that every convex subset Y of a normed space $(Z, \|\cdot\|)$ equipped with the metric induced from $(Z, \|\cdot\|)$ is an R-space, where the map r_ε is defined as $r_\varepsilon(y, z) = z + (\varepsilon/\|y - z\|) \cdot (y - z)$, if $\|y - z\| > \varepsilon$, and $r_\varepsilon(y, z) = y$, otherwise.

Theorem 4. *Let X be a topological space, (Y, d) be an R-space and $\mathcal{F}$ be a Δ-closed family of maps between X and Y. Then the family $\overline{\mathcal{F}}^{\mathrm{p}}$ is closed under uniform limits.*

Proof. Let $(r_n)_{n=1}^\infty$ be a sequence of continuous functions $r_n : Y \times Y \to Y$ such that

$$d(y, z) \le \frac{1}{2^n} \implies r_n(y, z) = y, \tag{3.3}$$

$$d(r_n(y, z), z) \le \frac{1}{2^n} \tag{3.4}$$

for all $y, z \in Y$ and $n \in \mathbb{N}$.

We consider a sequence $(f_n)_{n=1}^\infty$ of maps $f_n \in \overline{\mathcal{F}}^{\mathrm{p}}$ which is convergent to a map $f : X \to Y$ uniformly on X. Without loss of generality we may assume that

$$d(f_n(x), f_{n+1}(x)) \le \frac{1}{2^{n+1}}$$

for all $n \in \mathbb{N}$ and $x \in X$. For every n we take a sequence of maps $g_{n,m} : X \to Y$ such that

$$\lim_{m\to\infty} g_{n,m}(x) = f_n(x) \quad \forall x \in X,$$
$$g_{n,m} \in \mathcal{F} \quad \forall m, n \in \mathbb{N}.$$

For all $x \in X$ and $m \in \mathbb{N}$ let

$$\varphi_{1,m}(x) = g_{1,m}(x) \quad \text{and} \quad \varphi_{n,m}(x) = r_{n-1}(g_{n,m}(x), \varphi_{n-1,m}(x)) \quad \text{for } n > 1.$$

Then $\varphi_{n,m} \in \mathcal{F}$ for all n, m, since $\mathcal{F}$ is Δ-closed. Moreover, the inequality

$$d(\varphi_{n+1,m}(x), \varphi_{n,m}(x)) \le \frac{1}{2^n}$$

holds for all $n \in \mathbb{N}$ and $x \in X$.

We claim that the sequence $(\varphi_{n,m})_{m=1}^\infty$ converges to f_n pointwisely on X for every n. Indeed, fix $x \in X$ and show that for every $n \in \mathbb{N}$ there exists a number m_n such that $\varphi_{n,m}(x) = g_{n,m}(x)$ for all $m \ge m_n$. For $n = 1$ the statement is evident. Suppose that $n > 1$ and $\varphi_{n-1,m}(x) = g_{n-1,m}(x)$ for $m \ge m_{n-1}$. Since $(g_{n,m}(x))_{m=1}^\infty$ converges to $f_n(x)$ and $(g_{n-1,m}(x))_{m=1}^\infty$ converges to $f_{n-1}(x)$, there exists a number m_0 such that

$$d(g_{n,m}(x), f_n(x)) < \frac{1}{2^{n+1}} \quad \text{and} \quad d(g_{n-1,m}(x), f_{n-1}(x)) < \frac{1}{2^{n+1}}$$

for all $m \geq m_0$. Denote $m_n = \max\{m_0, m_{n-1}\}$ and notice that

$$d(g_{n,m}(x), \varphi_{n-1,m}(x)) = d(g_{n,m}(x), g_{n-1,m}(x)) \leq$$
$$\leq d(g_{n,m}(x), f_n(x)) + d(f_n(x), f_{n-1}(x)) + d(f_{n-1}(x), g_{n-1,m}(x)) \leq$$
$$\leq \frac{1}{2^{n+1}} + \frac{1}{2^n} + \frac{1}{2^{n+1}} = \frac{1}{2^{n-1}}$$

for all $m \geq m_n$. Consequently,

$$\varphi_{n,m}(x) = r_{n-1}(g_{n,m}(x), \varphi_{n-1,m}(x)) = g_{n,m}(x)$$

for all $m \geq m_n$.

Finally, we prove that $\lim_{m\to\infty} \varphi_{m,m}(x) = f(x)$. Fix $\varepsilon > 0$ and take $n_0 \in \mathbb{N}$ such that

$$\frac{1}{2^{n_0-1}} < \frac{\varepsilon}{3} \quad \text{and} \quad d(f_{n_0}(x), f(x)) \leq \frac{\varepsilon}{3}.$$

Since $\lim_{m\to\infty} \varphi_{n_0,m}(x) = f_{n_0}(x)$, there exists $k_0 > n_0$ such that

$$d(\varphi_{n_0,m}(x), f_{n_0}(x)) \leq \frac{\varepsilon}{3}$$

for all $m \geq k_0$. Then

$$d(\varphi_{m,m}(x), f(x)) \leq \sum_{i=n_0}^{m-1} d(\varphi_{i+1,m}(x), \varphi_{i,m}(x)) + d(\varphi_{n_0,m}(x), f_{n_0}(x)) + d(f_{n_0}(x), f(x)) \leq$$
$$\leq \sum_{i=n_0}^{m-1} \frac{1}{2^i} + \frac{\varepsilon}{3} + \frac{\varepsilon}{3} < \frac{1}{2^{n_0-1}} + \frac{\varepsilon}{3} + \frac{\varepsilon}{3} < \varepsilon$$

for all $m \geq k_0$. Therefore, $f \in \overline{\mathcal{F}}^{\mathrm{p}}$. □

We will use the following statement from [4] (see also [7, §31.VIII, Theorem 3] for metrizable domains).

Lemma 5. *Let X be a normal space and Y be a metric separable space. Then every map $f \in \mathrm{H}_1(X, Y)$ is a limit of uniformly convergent sequence of functions $f_n \in \mathrm{H}_1(X, Y)$ with countable discrete image $f_n(X)$ for every $n \in \mathbb{N}$. If, moreover, Y is completely bounded, we may assume that each function f_n takes finitely many values.*

4. Approximation of maps with path-connected range

Lemma 6. *Let P and X be as in Proposition 2 and let $f \in \mathrm{H}_1(X, \mathbb{R})$ has a discrete countable image $f(X)$. Then $f \in \overline{\mathrm{CF}(X)}^{\mathrm{p}}$.*

Proof. We consider a homeomorphism $\varphi : \mathbb{R} \to (0, 1)$ and the function $\varphi \circ f : X \to (0, 1)$ of the first Lebesgue class. By Lemma 5 we choose a sequence $(g_n)_{n=1}^{\infty}$ of functions from $\mathrm{H}_1^0(X, (0, 1))$ which is convergent to $\varphi \circ f$ uniformly on X. Since $g_n \in \overline{\mathrm{CF}(X, (0, 1))}^{\mathrm{p}}$ according to Corollary 3 for every $n \in \mathbb{N}$ and the space $Y = (0, 1)$ with the metric $d(x, y) = |x - y|$ is an R-space, $\varphi \circ f \in \overline{\mathrm{CF}(X, (0, 1))}^{\mathrm{p}}$ by Theorem 4. Hence, $f \in \overline{\mathrm{CF}(X)}^{\mathrm{p}}$. □

Proposition 7. *Let P and X be as in Proposition 2 and let Y be a path-connected space. Then every Lebesgue one function $f : X \to Y$ with countable discrete image $f(X)$ is a pointwise limit of a sequence of functions from $\mathrm{CF}(X, Y)$.*

Proof. Let $f \in \mathrm{H}_1(X, Y)$ and $f(X) = \{y_n : n \in \mathbb{N}\}$, $y_n \geq y_m$ for $n \geq m$. For every $n \in \mathbb{N}$ we take a continuous map $\gamma_n : [n, n + 1] \to Y$ such that $\gamma_n(n) = y_n$ and $\gamma_n(n + 1) = y_{n+1}$. Now we define a continuous map $\gamma : [1, +\infty) \to Y$ such that $\gamma|_{[n,n+1]} = \gamma_n$ for every $n \in \mathbb{N}$. Notice that every set $A_n = f^{-1}(y_n)$ is functionally F_σ in X and the family $(A_n : n \in \mathbb{N})$ forms a partition of X. We put $g(x) = n$ if $x \in A_n$ for some $n \in \mathbb{N}$. Then $g \in \mathrm{H}_1(X, [1, +\infty))$ and the space $g(X)$ is discrete

and countable. By Lemma 6 there exists a sequence of functions $g_m \in \mathrm{CF}(X)$ which is convergent to g pointwisely on X. Let us define functions $f_m \in \mathrm{CF}(X,Y)$ by the formula $f_m = \gamma \circ g_m$ for $m \in \mathbb{N}$. It remains to notice that for every $x \in X$ there exists $n \in \mathbb{N}$ such that $x \in A_n$, therefore, $\lim_{m\to\infty} f_m(x) = \gamma(g(x)) = \gamma(n) = y_n = f(x)$. □

Theorem 8. *Let P and X be as in Proposition 2 and (Y,d) be a path-connected separable R-space. Then every Lebesgue one function $f : X \to Y$ is a pointwise limit of a sequence of functions from $\mathrm{CF}(X,Y)$.*

Proof. Let $f \in \mathrm{H}_1(X,Y)$. According to Lemma 5 there exists a sequence of functions $f_n \in \mathrm{H}_1(X,Y)$ with discrete countable image which is convergent to f uniformly on X. Then $f_n \in \overline{\mathrm{CF}(X,Y)}^{\mathrm{p}}$ for every n by Proposition 7. Since the family $\mathrm{CF}(X,Y)$ is Δ-closed, $f \in \overline{\mathrm{CF}(X,Y)}^{\mathrm{p}}$ by Theorem 4. □

5. Semicontinuous functions

Theorem 9. *Let P and X be as in Proposition 2 and let $f : X \to [0,+\infty)$ be a lower semicontinuous function. Then f is a pointwise limit of an increasing sequence of functions $f_n \in \mathrm{CF}(X,[0,+\infty))$.*

Proof. Let $\mathbb{Q} = \{q_n : n \in \mathbb{N}\}$ and $B_n = (q_n, +\infty)$. Since f is lower semicontinuous, every set $G_n = f^{-1}(B_n)$ is open in X. Lemma 1 implies the existence of countable family $\mathcal{F} = (f_n : n \in \mathbb{N})$ of functions $f_n \in \mathrm{CF}(P,[0,1])$ such that every set G_n is open in $(X, d_{\mathcal{F}})$. Moreover, for every $a \geq 0$ there exists a sequence $(n_k)_{k=1}^{\infty}$ such that $f^{-1}(a,+\infty) = \bigcup_{k=1}^{\infty} G_{n_k}$. Then $f^{-1}(a,+\infty)$ is open in $(X, d_{\mathcal{F}})$. Therefore, $f : (X, d_{\mathcal{F}}) \to [0,+\infty)$ is lower semicontinuous.

For all $n \in \mathbb{N}$ and $x, y \in P$ we put

$$d_n(x,y) = \sum_{k=1}^{n} \frac{1}{2^k} |f_k(x) - f_k(y)|$$

and observe that

$$d_n(x,y) \leq d_{n+1}(x,y) \leq d_{\mathcal{F}}(x,y) \leq d_n(x,y) + \frac{1}{2^n}. \tag{5.1}$$

For all $n \in \mathbb{N}$ and $x \in X$ let

$$g_n(x) = \inf_{z\in X} (f(z) + n d_n(x,z)).$$

Then g_n depends on the first n coordinates, since for every fixed $z \in X$ the function $d_n(x,z)$ depends on on the first n coordinates. Moreover, $g_n(x) \leq f(x)$ for every $x \in X$.

Fix $n \in \mathbb{N}$ and notice that the inequalities

$$|g_n(x) - g_n(y)| \leq \sup_{z\in X} |n d_n(x,z) - n d_n(y,z)| \leq n d_n(x,y) \leq n d_{\mathcal{F}}(x,y)$$

for all $x, y \in X$ imply that g_n is uniformly continuous on $(X, d_{\mathcal{F}})$.

It follows from (5.1) that the sequence $(g_n)_{n=1}^{\infty}$ increases.

It remains to verify that $\lim_{n\to\infty} g_n(x) = f(x)$ for every $x \in X$. Fix $x \in X$ and $\varepsilon > 0$. Take a sequence $(z_n)_{n=1}^{\infty}$ of points from X with

$$f(z_n) + n d_n(x, z_n) < g_n(x) + \varepsilon.$$

Taking into account that $g_n(x) \leq f(x)$ for every n and $f(z_n) \geq 0$, we get $\lim_{n\to\infty} d_n(x,z_n) = 0$. Then $\lim_{n\to\infty} d_{\mathcal{F}}(x,z_n) = 0$. Since f is lower semicontinuous on $(X, d_{\mathcal{F}})$ at the point x, there exists a number k such that $f(z_n) \geq f(x) - \varepsilon$ for all $n \geq k$. Then $g_n(x) > f(z_n) - \varepsilon > f(x) - 2\varepsilon$ for all $n \geq k$. □

Corollary 10. *Let P and X be as in Proposition 2 and let $f : X \to \mathbb{R}$ be a lower semicontinuous function. Then f is a pointwise limit of an increasing sequence of functions from $\mathrm{CF}(X)$ if and only if f has a minorant from $\mathrm{CF}(X)$.*

Proof. Since the necessity is obvious, we will prove the sufficiency. Assume that $g \in \mathrm{CF}(X)$ is a minorant for f and put $h = f - g$. Applying Theorem 9 for the function $h : X \to [0, +\infty)$, we obtain the increasing sequence of functions $h_n \in \mathrm{CF}(X, [0, +\infty))$ which is convergent to h pointwisely on X. It remains to put $f_n = h_n + g$ for every $n \in \mathbb{N}$. □

6. Pseudocompact case

We say that a sequence $(f_n)_{n=1}^{\infty}$ of maps $f_n : X \to Y$ between topological spaces X and Y is *stably convergent to a map $f : X \to Y$ on X*, if for every $x \in X$ there exists $k \in \mathbb{N}$ such that $f_n(x) = f(x)$ for all $n \geq k$.

Lemma 11. *Let $f : X \to Y$ be a map between a topological space X and a metric space (Y, d_Y), $(f_n)_{n=1}^{\infty}$ be a sequence of maps $f_n : X \to Y$ which is convergent to f stably on X and let $(f_{n,k})_{k=1}^{\infty}$ be a sequence of maps $f_{n,k} : X \to Y$ which is convergent to f_n uniformly on X for every $n \in \mathbb{N}$. Then $f \in \overline{\{f_{n,k} : n, k \in \mathbb{N}\}}^{\mathrm{p}}$.*

Proof. Without loss of generality we may assume that

$$d_Y(f_{n,k}(x), f_n(x)) < \frac{1}{2^k},$$
$$d_Y(f_{n,k+1}(x), f_{n,k}(x)) < \frac{1}{2^k}$$

for all $x \in X$ and $n, k \in \mathbb{N}$.

Fix $x \in X$ and $\varepsilon > 0$. We choose $m \in \mathbb{N}$ such that

$$\frac{1}{2^{m-3}} < \varepsilon \quad \text{and} \quad f_n(x) = f(x) \quad \forall n \geq m.$$

Then we get

$$d_Y(f_{n,n}(x), f(x)) \leq \sum_{k=m}^{n-1} d_Y(f_{n,k+1}(x), f_{n,k}(x)) + d_Y(f_{n,m}(x), f_n(x)) <$$
$$< \sum_{k=m}^{n-1} \frac{1}{2^k} + \frac{1}{2^m} < \frac{1}{2^{m-2}} + \frac{1}{2^m} < \frac{1}{2^{m-3}} < \varepsilon$$

for all $n \geq m$. Therefore, $\lim\limits_{n\to\infty} f_{n,n}(x) = f(x)$. □

Lemma 12. *Let $P = \prod_{n=1}^{\infty} X_n$ be a pseudocompact space and (Y, d_Y) be a path-connected metric space. Then every Baire one function $f : X \to Y$ with countable discrete image $f(X)$ is a pointwise limit of a sequence of functions from $\mathrm{CF}(X, Y)$.*

Proof. It follows from [5, Lemma 4] that there exists a sequence $(f_n)_{n=1}^{\infty}$ of continuous functions $f_n : P \to Y$ which is stably convergent to f on P. By [1, Theorem 1] every function f_n is the uniform limit of a sequence from $\mathrm{CF}(P, Y)$. It remains to apply Lemma 11. □

Theorem 13. *Let $P = \prod_{n=1}^{\infty} X_n$ be a pseudocompact space and (Y, d_Y) be a path-connected separable metric R-space. Then every Baire one function $f : P \to Y$ is a pointwise limit of a sequence of functions from $\mathrm{CF}(P, Y)$.*

Proof. Let $f \in \mathrm{B}_1(P, Y)$. It follows from [4, Lemma 3.3] and [5, Lemma 4] that there exists a sequence of functions $f_n \in \mathrm{B}_1(P, Y)$ with discrete countable image $f(P)$ which is uniformly convergent to f. Then $f_n \in \overline{\mathrm{CF}(P, Y)}^{\mathrm{p}}$ for every n by Lemma 12. Since the family $\mathrm{CF}(P, Y)$ is Δ-closed, $f \in \overline{\mathrm{CF}(P, Y)}^{\mathrm{p}}$ by Theorem 4. □

The authors do not know an answer to the following question.

Question 3. *Let $X \subseteq \prod_{n=1}^{\infty} X_n$ be a pseudocompact subspace of a product of completely regular spaces X_n and $f : X \to \mathbb{R}$ be a Baire one function. Does there exist a sequence of functions from $\mathrm{CF}(X)$ which is pointwisely convergent to f on X?*

7. Example

Denote by $\mathbf{C}$ the Cantor set $\mathbf{C} \subseteq [0,1]$. For every $n \in \mathbb{N}$ and $k \in \{1, \dots, n\}$ we denote by $p_{k,n}$ the natural projection $p_{k,n} : \mathbf{C}^n \to \mathbf{C}$, $p_{k.n}(x_1, \dots x_n) = x_k$. We put

$$\mathrm{r}_n(A) = \bigcup_{k=1}^{n} p_{k,n}^{-1}(A)$$

for every $n \in \mathbb{N}$ and for an infinite set $A \subseteq \mathbf{C}$. Following to [11, Fact 3, p. 124], for every $n \in \mathbb{N}$ and $P \subseteq \mathbf{C}^n$ we put

$$|P|_n = \min\{|A| : A \subseteq \mathbf{C} \text{ and } P \subseteq \mathrm{r}_n(A)\}.$$

According to [11, Fact 4, p. 125], we have that if $|F|_n > \aleph_0$ for some $n \in \mathbb{N}$ and a closed in $\mathbf{C}^n$ set F then $|F|_n = \mathfrak{c}$.

Lemma 14. *There exists a sequence $(A_n)_{n=1}^{\infty}$ of sets $A_n \subseteq \mathbf{C}$ such that the following conditions hold:*

(1) $\mathbf{C} = \bigsqcup_{n=1}^{\infty} A_n$;
(2) $(A_n)^m \cap F \neq \emptyset$ *for all* $n, m \in \mathbb{N}$ *and a closed set* F *in* $\mathbf{C}^m$ *with* $|F|_m = \mathfrak{c}$.

Proof. For every $m \in \mathbb{N}$ we denote by $\mathcal{F}_m$ the system of all closed sets $F \subseteq \mathbf{C}^m$ with $|F|_m = \mathfrak{c}$. Clearly, $|\mathcal{F}_m| = \mathfrak{c}$ for every $m \in \mathbb{N}$. Let $\bigcup_{m=1}^{\infty} \mathcal{F}_m = \{F_\xi : 1 \leq \xi < \mathfrak{c}\}$. For every $\xi < \mathfrak{c}$ we denote by m_ξ an integer $m_\xi \in \mathbb{N}$ such that $F_\xi \subseteq \mathbf{C}^{m_\xi}$. By transfinite induction on ξ we construct a family $((A_{\xi,n})_{n=1}^{\infty} : 1 \leq \xi < \mathfrak{c})$ of sequences $(A_{\xi,n})_{n=1}^{\infty}$ of finite sets $A_{\xi,n} \subseteq \mathbf{C}$ which satisfies the following conditions:

(a) $A_{\xi,n} \cap A_{\theta,m} = \emptyset$ for $(\xi, n) \neq (\theta, m)$;
(b) $(A_{\xi,n})^{m_\xi} \cap F_\xi \neq \emptyset$ for every $\xi < \mathfrak{c}$ and $n \in \mathbb{N}$.

Assume that $1 \leq \alpha < \mathfrak{c}$ and the family $((A_{\xi,n})_{n=1}^{\infty} : 1 \leq \xi < \alpha)$ is constructed. We put

$$B_\alpha = \bigcup_{1 \leq \xi < \alpha} \bigcup_{n=1}^{\infty} A_{\xi,n}.$$

Clearly, $|B_\alpha| < \mathfrak{c}$. Since $\mathrm{r}_{m_\alpha}(F_\alpha) = \mathfrak{c}$, we can construct easily by the induction on n a sequence $(A_{\alpha,n})_{n=1}^{\infty}$ of disjoint finite sets $A_{\alpha,n} \subseteq \mathbf{C} \setminus B_\alpha$ such that $(A_{\alpha,n})^{m_\alpha} \cap F_\alpha \neq \emptyset$ for every $n \in \mathbb{N}$. It remains to put

$$A_n = \bigcup_{1 \leq \xi < \alpha} A_{\xi,n}$$

for every $n \geq 2$ and

$$A_1 = \mathbf{C} \setminus \left(\bigcup_{n \geq 2} A_n \right).$$

□

Lemma 15. *There exists a function $g : \mathbf{C} \to \mathbb{R}$ of the second Baire class such that for every countable set $A \subseteq \mathbf{C}$ the restriction $g|_{\mathbf{C} \setminus A}$ is not a function of the first Baire class.*

Proof. Let μ be the Lebesgue measure on $[0,1]$. We prove the statement for a function $g : X \to \mathbb{R}$ defined on a Cantor type closed set $X \subseteq [0,1]$ which is homeomorphic to $\mathbf{C}$ and $\mu(G) > 0$ for every open set $G \subseteq X$.

We choose an F_σ-set B of the first category in X such that $\mu(B) = \mu(X)$ and put $g = \chi|_B$. Since B is F_σ, the function g belongs to the second Baire class. Moreover, for every countable set

$A \subseteq X$ the function $g|_{X\setminus A}$ is everywhere discontinuous. Therefore, the restriction $g|_{X\setminus A}$ is not a Baire one function, because the space $X \setminus A$ is Baire. □

Theorem 16. *There exist a sequence $(X_n)_{n=1}^{\infty}$ of Lindelöf spaces X_n and a function $f : \prod_{n=1}^{\infty} X_n \to \mathbb{R}$ of the first Baire class such that*

(i) *for every $n \in \mathbb{N}$ the product $Y_n = \prod_{k=1}^{n} X_k$ is Lindelöf;*
(ii) *f is not a pointwise limit of any sequence $(f_n)_{n=1}^{\infty}$ of functions from $\mathrm{CF}(\prod_{n=1}^{\infty} X_n)$.*

Proof. Let $(A_n)_{n=1}^{\infty}$ be a sequence of sets $A_n \subseteq \mathbf{C}$ which satisfy conditions (1) and (2) of Lemma 14. Let τ_0 be the usual topology of the Cantor set $\mathbf{C}$. For every $n \in \mathbb{N}$ we denote by τ_n the topology on $\mathbf{C}$ with the base

$$\tau_0 \cup \{\{x\} : x \in A_n\}.$$

Put $X_n = (\mathbf{C}, \tau_n)$ for every $n \in \mathbb{N}$.

(i). Fix $n \in \mathbb{N}$ and put $B_n = \bigcup_{k=1}^{n} A_k$. Consider the space $Z = (\mathbf{C}, \sigma)$, where σ is a topology on $\mathbf{C}$ with the base

$$\tau_0 \cup \{\{x\} : x \in B_n\}.$$

According to [11, Fact 6, p. 126], the space Z^n is Lindelöf. Since the topology of the product Y_n is weaker than the topology of Z^n, Y_n is Lindelöf too.

(ii). We start with auxiliary properties of functions on the space $X = \prod_{n=1}^{\infty} X_n$.

`Claim A`. *For every function $v : \mathbf{C} \to \mathbb{R}$ of the first Baire class there exists a continuous function $u : X \to \mathbb{R}$ such that $u(a, a, \dots) = v(a)$ for every $a \in \mathbf{C}$.*

`Proof of Claim A.` Choose a sequence $(v_n)_{n=1}^{\infty}$ of continuous functions $v_n : \mathbf{C} \to \mathbb{R}$ which converges to v pointwisely on $\mathbf{C}$. Using the uniform continuity of the functions v_n, we choose a strictly decreasing sequence of reals $\delta_n > 0$ such that $|v_n(x') - v_n(x'')| < \frac{1}{n}$ for every $n \in \mathbb{N}$ and $x', x'' \in \mathbf{C}$ with $|x' - x''| < \delta_n$. For every $n \in \mathbb{N}$ we put

$$G_n = \{(x_k)_{k=1}^{\infty} \in X : \max\{|x_i - x_j| : 1 \le i, j \le n\} < \delta_n\}$$

and

$$F_n = \{(x_k)_{k=1}^{\infty} \in X : \max\{|x_i - x_j| : 1 \le i, j \le n\} \le \delta_n\}.$$

It is evident that every set G_n is open and every set F_n is closed in $\mathbf{C}^{\omega}$. We put $v_0(x) = 0$ for every $x \in \mathbf{C}$ and

$$\Delta = \{(x, x, \dots) : x \in \mathbf{C}\}.$$

For every $n \in \mathbb{N}$ we choose a continuous function $\varphi_n : \mathbf{C}^{\omega} \to [0, 1]$ such that $\varphi_n(X \setminus G_n) \subseteq \{1\}$ and $\varphi_n(F_{n+1}) \subseteq \{0\}$ and consider the function $u : X \to \mathbb{R}$,

$$u(x) = \begin{cases} v_0(x), & x \in X \setminus F_1 \\ \varphi_n(x)v_{n-1}(x_1) + (1 - \varphi_n(x)v_n(x_1), & x = (x_1, x_2, \dots) \in F_n \setminus F_{n+1} \\ v(a), & x = (a, a, \dots) \in \Delta. \end{cases}$$

It easy to see that the function u is continuous at every point $x \in X \setminus \Delta$. It remains to verify the continuity of u at every point $x \in \Delta$. Fix $a \in \mathbf{C}$ and $\varepsilon > 0$. We choose $m \in \mathbb{N}$ such that $a \in A_m$ and let $n_0 \ge \max\{m, \frac{2}{\varepsilon} + 1\}$ be such that $|v_n(a) - v(a)| < \frac{\varepsilon}{2}$ for all $n \ge n_0 - 1$. Now we consider the neighborhood

$$U = \{(x_n)_{n=1}^{\infty} \in X : x_m = a \text{ and } \max\{|x_i - x_j| : 1 \le i, j \le n_0\} \le \delta_{n_0}\}$$

of the point $x_0 = (a, a, \dots)$ in X. Fix $x = (x_n)_{n=1}^{\infty} \in U \setminus \{x_0\}$ and take $k \in \mathbb{N}$ such that $x \in F_k \setminus F_{k+1}$. Notice that $k \ge n_0$, because $U \subseteq F_{n_0}$. Therefore,

$$|v_k(a) - v(a)| < \frac{\varepsilon}{2} \quad \text{and} \quad |v_{k-1}(a) - v(a)| < \frac{\varepsilon}{2}.$$

Since $x_m = a$, $|x_1 - a| < \delta_k$ and, moreover $|x_1 - a| < \delta_{k-1}$. Therefore,

$$|v_k(x_1) - v_k(a)| < \tfrac{1}{k} \le \tfrac{1}{n_0} \le \tfrac{\varepsilon}{2}$$

and

$$|v_{k-1}(x_1) - v_{k-1}(a)| < \tfrac{1}{k-1} \le \tfrac{1}{n_0-1} \le \tfrac{\varepsilon}{2}.$$

Thus,

$$|v_k(x_1) - v(a)| < \varepsilon \quad \text{and} \quad |v_{k-1}(a) - v(a)| < \varepsilon.$$

Now we have

$$|u(x) - v(a)| = |\varphi_k(x)(v_{k-1}(x_1) - v(a)) + (1 - \varphi_k(x)(v_k(x_1) - v(a)| \le$$

$$\varphi_k(x)|v_{k-1}(x_1) - v(a)| + (1 - \varphi_k(x)|v_k(x_1) - v(a)| < \varphi_k(x)\varepsilon + (1 - \varphi_k(x)\varepsilon = \varepsilon.$$

`Claim B`. *For every function $w : \mathbf{C} \to \mathbb{R}$ of the second Baire class the function $h : X \to \mathbb{R}$*

$$h(x) = \begin{cases} 0, & x \in X \setminus \Delta \\ w(a), & x = (a, a, a, \dots) \in \Delta \end{cases}$$

is a function of the first Baire class.

`Proof of Claim B.` We choose a sequence $(w_n)_{n=1}^{\infty}$ of Baire one functions $w_n : \mathbf{C} \to \mathbb{R}$ which converges to w pointwisely on $\mathbf{C}$. By Claim A we can construct a sequence $(h_n)_{n=1}^{\infty}$ of continuous functions $h_n : X \to \mathbb{R}$ such that $h_n(a, a, \dots) = w_n(a)$ for all $a \in X$ and $h_n(x) = 0$ for all $n \in \mathbb{N}$ and $x \in X \setminus C_n$, where

$$C_n = \{(x_k)_{k=1}^{\infty} \in X : \max\{|x_i - x_j| : 1 \le i, j \le n\} \le \tfrac{1}{n}\}.$$

Clearly, the sequence $(h_n)_{n=1}^{\infty}$ converges to h pointwisely on X.

Now we prove the condition (ii). Lemma 15 implies the existence of a function $g : \mathbf{C} \to \mathbb{R}$ of the second Baire class such that for every countable set $A \subseteq \mathbf{C}$ the restriction $g|_{\mathbf{C}\setminus A}$ does not belong to the first Baire class. According to Claim B, the function $f : X \to \mathbb{R}$

$$f(x) = \begin{cases} 0, & x \in X \setminus \Delta \\ g(a), & x = (a, a, a, \dots) \in \Delta \end{cases}$$

is a function of the first Baire class.

Now we show that (ii) holds. Suppose to the contrary that f is the pointwise limit of a sequence of functions $f_n \in \mathrm{CF}(X)$. Without loss of generality we may assume that for every $n \in \mathbb{N}$ the function f_n depends on the first n coordinates. For every $n \in \mathbb{N}$ and $z \in \mathbf{C}$ we put

$$g_n(z) = f_n(z, z, \dots).$$

Fix $n \in \mathbb{N}$. Since f_n depends on the first n coordinates, the function $g_n : \mathbf{C} \to \mathbb{R}$ is continuous at every point $z \in \mathbf{C} \setminus (\bigcup_{k=1}^{n} A_k)$. It follows from condition (2) of Lemma 14 that every closed set $F \subseteq \bigcup_{k=1}^{n} A_k$ is at most countable. Therefore, an F_σ-set D_n of all discontinuity points of f_n is at most countable too.

Hence, the set $D = \bigcup_{n=1}^{\infty} D_n$ is at most countable, every function g_n is continuous at each point $z \in \mathbf{C} \setminus D$ and the sequence $(g_n)_{n=1}^{\infty}$ converges to g pointwisely on $\mathbf{C}$, which contradicts to the choice of g. □

Remark 2. 1) *It easy to see that for a product $X = \prod_{n=1}^{\infty} X_n$ of a sequence $(X_n)_{n=1}^{\infty}$ of topological spaces X_n and for a continuous map $f : X \to Y$ with values in a topological space Y there exists a sequence $(f_n)_{n=1}^{\infty}$ of functions $f_n \in \mathrm{CF}(X, Y)$ which converges to f pointwisely on X.*

2) *In fact it was proved in Theorem 8 that for appropriate spaces $X \subseteq \prod_{n=1}^{\infty} X_n$ and Y for every mapping $f : X \to Y$ of the first Baire class there exists a sequence of functions from $\mathrm{CF}(\prod_{n=1}^{\infty} X_n, Y)$ such that the sequence of the restrictions $f_n|_X$ converges to f pointwisely on X.*

3) *For the sequence $(X_n)_{n=1}^{\infty}$ of topological spaces X_n from Theorem 16, for the subspace $X = \Delta$ and $f \in \mathrm{CF}(X)$ with $f(x, x, \dots) = g(x)$, there does not exist a sequence of functions $f_n : \prod_{n=1}^{\infty} X_n \to \mathbb{R}$ such that the sequence of the restrictions $f_n|_X$ converges to f pointwisely.*

References

[1] V. Bykov, *On Baire class one functions on a product space*, Topol. Appl. **199** (2016) 55–62.
[2] R. Engelking, *On functions defined on Cartesian products*, Fund. Math. **59** (1966) 221-231.
[3] P. Hajek, *Smooth norms that depends locally on finitely many coordinates*, Proc. Amer. Math. Soc. **123 (12)** (1995) 3817–3821.
[4] O. Karlova, *The first functional Lebesgue class and Baire classification of separately continuous mappings*, Nauk. Visn. Chernivtsi University. Matematyka, **191-192** (2004), 52–60 (in Ukrainian).
[5] O. Karlova, *Baire classification of maps with values in subsets of finitely dimensional spaces*, Nauk. Visn. Chernivtsi University. Matematyka, **239** (2005), 59–65 (in Ukrainian).
[6] O. Karlova, *On Baire classification of mappings with values in connected spaces*, Eur. J. Math. (2016) DOI:10.1007/s40879-015-0076-y.
[7] K. Kuratowski, *Topology. Volume 1*, Academic Press, New York, 1966.
[8] S. Mazur, *On continuous mappings on Cartesian products*, Fund. Math. **39** (1952) 229-238.
[9] V. Mykhaylyuk, *The Namioka property of KC-functions andKempisty spaces*, Topol. Appl. **153** (2006) 2455–2461.
[10] N. Noble, M. Ulmer, *Factoring functions on Cartesian products*, Trans. Amer. Math. Soc. **163** (1072) 329-339.
[11] V. Tkachuk, *A C_p-Theory Problem Book: Topological and Function Spaces (Problem Books in Mathematics)*, Springer New York Dordrecht Heidelberg London (2011), 485.

Department of Mathematical Analysis, Faculty of Mathematics and Informatics, Yurii Fedkovych Chernivtsi National University, 58000 Kotsyubyns'koho str., 2, Chernivtsi, Ukraine